



PCT/GB 04/4481



INVESTOR IN PEOPLE

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

REC'D 04 NOV 2004

WIPO

PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

Signed

Andrew Gersey

Dated

13 September 2004

**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

BEST AVAILABLE COPY

The
Patent
Office

1/77

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

THE PATENT OFFICE

C

24 OCT 2003

NEWPORT

The Patent Office

Cardiff Road
Newport
Gwent NP9 1RH

1. Your reference

101224

24 OCT 2003

24OCT03 E846958-4 D02934

P01/7700 0.00-0324791.3

2. Patent application number

(The Patent Office will fill in this part)

0324791.3

3. Full name, address and postcode of the or of each applicant (underline all surnames)

~~AstraZeneca AB~~

~~SE 151 85 Sodertalje~~

~~Sweden~~

ASTRAZENECA UK LIMITED
15 STANHOPE GATE
LONDON
W1K 1LN

Sweden

Patents ADP number (if you know it)

7898467002

If the applicant is a corporate body, give the country/state of its incorporation

4. Title of the invention

CHEMICAL PROCESS

5. Name of your agent (if you have one)

Kevin Bill

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

AstraZeneca
Global Intellectual Property
PO Box 272
Mereside, Alderley Park
Macclesfield, Cheshire SK10 4TG

Patents ADP number (if you know it)

08179707001

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application number
(if you know it)

Date of filing
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

a) any applicant named in part 3 is not an inventor, or

b) there is an inventor who is not named as an applicant, or

c) any named applicant is a corporate body.

See note (d))

Patents Form 1/77

Enter the number of sheets for any of the following items you are filing with this form.
Do not count copies of the same document

Continuation sheets of this form

Description 27

Claim(s) 04

Abstract

Drawing(s)

0. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77)

Request for substantive examination (Patents Form 10/77)

Any other documents (please specify)

11.

I/We request the grant of a patent on the basis of this application.

Signature

Date

Authorised Signatory

23/10/2003

12. Name and daytime telephone number of person to contact in the United Kingdom

Helen Dixon - 01625 517301

Warning


After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

CHEMICAL PROCESS


This invention concerns improvements to a chemical process, particularly a chemical process for manufacture of (*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoic acid calcium salt (1) (illustrated below), which is useful for the production of a pharmaceutical useful in the treatment of, inter alia, hypercholesterolemia, hyperlipoproteinemia and atherosclerosis. This invention also concerns crystalline intermediates useful in the chemical process.



1


The sodium salt (3) and calcium salt (1) of the compound (*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoic acid (hereinafter referred to as the 'Agent') were disclosed in European Patent 0521471.

This patent also describes a process for the synthesis of the calcium salt (1), via the dihydroxy ester (2) and the sodium salt (3), as shown in Scheme 1 below. The calcium salt thus formed is then collected and dried and may be processed further as required.



Scheme 1

Our International Patent Application WO 00/49014 describes an alternative route to the calcium salt (1), also via the sodium salt (3), from the compound tert-butyl (E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]vinyl}(4R,6S)-2,2-dimethyl[1,3]dioxan-4-yl)acetate (BEM) (4), which is exemplified as shown in Scheme 2 below:



Scheme 2

As described in WO 00/49014, the transformation from BEM (4) to the calcium salt (1) may be carried out via the methylamine salt (5) as shown in Scheme 2. Isolation of this intermediate crystalline methylamine salt allows purification by recrystallisation before final formation of the (amorphous) calcium salt. However formation of the methylamine salt introduces an extra step into the process, which is generally undesirable for manufacture (for example because of additional cost and the potential for introduction of additional impurities).

The transformation from BEM (4) to the sodium salt (3) in Scheme 2 takes place in two steps as shown below in Scheme 3.


Scheme 3

5 salt (3) of the parent carboxylic acid. The intermediate compound BED (6) is not isolated in the process described in WO 00/49014. The analogous methyl ester (2) was described in European Patent 0521471 as a syrup and therefore, by analogy, isolation of BED (6) would not be expected to provide any advantages to the process.

10 are generally crystalline compounds which may advantageously be isolated and recrystallised, thus removing the need for isolation of an intermediate salt such as the methylamine salt in order for purification to be carried out.


15 yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoic acid, comprising:

a) acid hydrolysis of an acetal protecting group in a compound of the formula (7)




7

wherein A is an acetal or ketal protecting group and R is (1-6C)alkyl, and isolation of the resulting crystalline compound of the formula (8);



8

- 5 b) optional recrystallisation of the compound of the formula (8);
 c) hydrolysis of the ester group in the compound (8) to give a dihydroxy carboxylate derivative (9) (wherein M is hydrogen or a metal counterion other than calcium) or a compound of the formula (1); and



9

- 10 d) where necessary, conversion of a compound of the formula (9) into a compound of the formula (1).

In a further aspect of the invention there is provided crystalline compounds of the formula (8).

- Furthermore, we have found that some compounds of the formula (7), which are
 15 analogues of BEM (4) are also crystalline and are themselves useful intermediates which may

be recrystallised if necessary to improve the quality of the material. These crystalline compounds are novel and each independently provides a further aspect of the invention.

Thus in a further aspect of the invention is provided crystalline methyl (E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]vinyl})(4R,6S)-
5 2,2-dimethyl[1,3]dioxan-4-yl)acetate having an X-ray powder diffraction pattern with peaks at 2-theta = 9.5, 13.6 and 17.5.

In a further aspect of the invention is provided crystalline methyl (E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]vinyl})(4R,6S)-
10 2,2-dimethyl[1,3]dioxan-4-yl)acetate having an X-ray powder diffraction pattern with peaks at 2-theta = 5.7, 9.5, 13.6, 17.5, 19.9 and 22.4.

In a further aspect of the invention is provided crystalline methyl (E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]vinyl})(4R,6S)-
2,2-dimethyl[1,3]dioxan-4-yl)acetate having an X-ray powder diffraction pattern with peaks at 2-theta = 5.7, 8.7, 9.5, 13.6, 17.5, 19.0, 19.9, 20.8, 21.8 and 22.4.

15 In a further aspect of the invention is provided crystalline methyl (E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]vinyl})(4R,6S)-
2,2-dimethyl[1,3]dioxan-4-yl)acetate having an X-ray powder diffraction pattern substantially as shown in Figure 1.

In a further aspect of the invention is provided crystalline ethyl (E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]vinyl})(4R,6S)-
20 2,2-dimethyl[1,3]dioxan-4-yl)acetate having an X-ray powder diffraction pattern with peaks at 2-theta = 15.9, 18.4 and 19.5.

In a further aspect of the invention is provided crystalline ethyl (E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]vinyl})(4R,6S)-
25 2,2-dimethyl[1,3]dioxan-4-yl)acetate having an X-ray powder diffraction pattern with peaks at 2-theta = 15.9, 18.4, 19.5, 23.0, 24.3 and 25.0.

In a further aspect of the invention is provided crystalline ethyl (E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]vinyl})(4R,6S)-
30 2,2-dimethyl[1,3]dioxan-4-yl)acetate having an X-ray powder diffraction pattern with peaks at 2-theta = 5.9, 8.0, 12.2, 15.9, 18.4, 19.5, 19.7, 23.0, 24.3 and 25.0.

In a further aspect of the invention is provided crystalline ethyl (E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]vinyl})(4R,6S)-

2,2-dimethyl[1,3]dioxan-4-yl)acetate having an X-ray powder diffraction pattern substantially as shown in Figure 2.

In a further aspect of the invention is provided crystalline iso-propyl (E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]vinyl}(4R,6S)-2,2-dimethyl[1,3]dioxan-4-yl)acetate having an X-ray powder diffraction pattern with peaks at 2-theta = 7.8, 11.6 and 15.5.

In a further aspect of the invention is provided crystalline iso-propyl (E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]vinyl}(4R,6S)-2,2-dimethyl[1,3]dioxan-4-yl)acetate having an X-ray powder diffraction pattern with peaks at 2-theta = 6.9, 7.0, 7.8, 8.7, 11.6 and 15.5.

In a further aspect of the invention is provided crystalline iso-propyl (E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]vinyl}(4R,6S)-2,2-dimethyl[1,3]dioxan-4-yl)acetate having an X-ray powder diffraction pattern with peaks at 2-theta = 6.9, 7.0, 7.8, 8.7, 10.4, 11.6, 13.0, 14.7, 15.5 and 20.2.

In a further aspect of the invention is provided crystalline iso-propyl (E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]vinyl}(4R,6S)-2,2-dimethyl[1,3]dioxan-4-yl)acetate having an X-ray powder diffraction pattern substantially as shown in Figure 3.

In a further aspect of the invention is provided crystalline n-hexyl (E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]vinyl}(4R,6S)-2,2-dimethyl[1,3]dioxan-4-yl)acetate having an X-ray powder diffraction pattern with peaks at 2-theta = 5.3, 7.1 and 18.9.

In a further aspect of the invention is provided crystalline n-hexyl (E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]vinyl}(4R,6S)-2,2-dimethyl[1,3]dioxan-4-yl)acetate having an X-ray powder diffraction pattern with peaks at 2-theta = 5.3, 7.1, 14.2, 14.8, 18.9 and 21.4.

In a further aspect of the invention is provided crystalline n-hexyl (E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]vinyl}(4R,6S)-2,2-dimethyl[1,3]dioxan-4-yl)acetate having an X-ray powder diffraction pattern with peaks at 2-theta = 5.3, 7.1, 11.4, 14.2, 14.8, 18.9, 20.1, 20.4 and 21.4.

In a further aspect of the invention is provided crystalline n-hexyl (E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]vinyl}(4R,6S)-

2,2-dimethyl[1,3]dioxan-4-yl)acetate having an X-ray powder diffraction pattern substantially as shown in Figure 4.

In a further aspect of the invention is provided crystalline ethyl-(*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoate having an X-ray powder diffraction pattern with peaks at 2-theta = 8.1, 11.3 and 19.9.

In a further aspect of the invention is provided crystalline ethyl-(*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoate having an X-ray powder diffraction pattern with peaks at 2-theta = 8.1, 11.3, 12.4, 19.9, 21.0 and 22.1.

In a further aspect of the invention is provided crystalline ethyl-(*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoate having an X-ray powder diffraction pattern with peaks at 2-theta = 4.3, 8.1, 11.3, 12.4, 15.1, 19.9, 21.0, 21.7, 22.1 and 23.5.

In a further aspect of the invention is provided crystalline ethyl-(*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoate having an X-ray powder diffraction pattern substantially as shown in Figure 5.

In a further aspect of the invention is provided crystalline iso-propyl-(*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoate having an X-ray powder diffraction pattern with peaks at 2-theta = 9.8, 17.3 and 21.1.

In a further aspect of the invention is provided crystalline iso-propyl-(*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoate having an X-ray powder diffraction pattern with peaks at 2-theta = 9.8, 12.2, 17.3, 19.6, 20.1 and 21.1.

In a further aspect of the invention is provided crystalline iso-propyl-(*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoate having an X-ray powder diffraction pattern with peaks at 2-theta = 9.8, 12.2, 13.6, 17.3, 18.5, 19.6, 20.1, 21.1, 22.4 and 23.3.

In a further aspect of the invention is provided crystalline iso-propyl-(*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-

dihydroxyhept-6-enoate having an X-ray powder diffraction pattern substantially as shown in Figure 6.

In a further aspect of the invention is provided crystalline tert-butyl-(*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoate having an X-ray powder diffraction pattern with peaks at 2-theta = 18.2, 19.9 and 20.8.

In a further aspect of the invention is provided crystalline tert-butyl-(*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoate having an X-ray powder diffraction pattern with peaks at 2-theta = 9.8, 18.2, 19.9, 20.6, 20.8 and 26.3.

In a further aspect of the invention is provided crystalline tert-butyl-(*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoate having an X-ray powder diffraction pattern with peaks at 2-theta = 9.8, 17.4, 18.2, 19.4, 19.9, 20.6, 20.8, 22.1, 25.1 and 26.3.

In a further aspect of the invention is provided crystalline tert-butyl-(*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoate having an X-ray powder diffraction pattern substantially as shown in Figure 7.


A crystalline polymorph of BEM (4) is provided as a further aspect of the invention.

The X-ray powder diffraction spectra were determined by mounting a sample of the crystalline form on Siemens single silicon crystal (SSC) wafer mounts and spreading out the sample into a thin layer with the aid of a microscope slide. The sample was spun at 30 revolutions per minute (to improve counting statistics) and irradiated with X-rays generated by a copper long-fine focus tube operated at 40kV and 40mA with a wavelength of 1.5406 angstroms. The collimated x-ray-source was passed through an automatic variable divergence slit set at V20 (20mm path length) and the reflected radiation directed through a 2mm antiscatter slit and a 0.2mm detector slit. The sample was exposed for 4 seconds per 0.02 degree 2-theta increment (continuous scan mode) over the range 2 degrees to 40 degrees 2-theta in theta-theta mode. The running time was 2 hours 6 minutes and 40 seconds. The instrument was equipped with a scintillation counter as detector. Control and data capture was by means of a DECpc LPv 433sx personal computer running with Diffrac AT (Socabim) software.

It will be understood that the 2-theta values of an X-ray powder diffraction pattern may vary slightly from one machine to another or from one sample to another, and so the values quoted are not to be construed as absolute. It will also be understood that the relative intensities of peaks may vary according to the orientation of the sample under test so that the intensities in the XRD traces included herein are illustrative and not intended to be used for absolute comparison.

The crystalline forms obtained according to the present invention are substantially free from other crystal and non-crystal forms of each compound of the formula 7 or 8. The term "substantially free from other crystal and non-crystal forms" shall be understood to mean that the desired crystal form contains less than 50%, preferably less than 10%, more preferably less than 5% of any other form of the compound.

Suitable values for the acetal protecting group A are as described in EP0319847. A preferred value for A is iso-propylidene such that the compound of the formula (7) is a compound of the formula (7a).



7a


Suitably R is selected from methyl, ethyl, propyl, iso-propyl, butyl, tert-butyl, pentyl and n-hexyl.

Preferably R is selected from methyl, ethyl, iso-propyl, tert-butyl and n-hexyl.

More preferably R is selected from ethyl, iso-propyl and tert-butyl.


In a further aspect of the invention there is provided a process for the manufacture of the calcium salt of (*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoic acid, comprising:

a) acid hydrolysis of an acetal protecting group in a compound of the formula (7a)




7a

wherein R is ethyl, iso-propyl or tert-butyl, and isolation of the resulting crystalline compound of the formula (8);



8

- 5 b) optional recrystallisation of the compound of the formula (8);
 c) hydrolysis of the ester group in the compound (8) to give a dihydroxy carboxylate derivative (9) (wherein M is hydrogen or a metal counterion other than calcium) or a compound of the formula (1); and




9

- 10 d) where necessary, conversion of a compound of the formula (9) into a compound of the formula (1).


In a further aspect of the invention there is provided a process for the manufacture of the calcium salt of (*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoic acid,

- 15 comprising:



8

a) hydrolysis of the ester group in a crystalline compound of the formula (8) (where R is as hereinbefore defined) to give a dihydroxy carboxylate derivative (9) (wherein M is hydrogen or a metal counterion other than calcium) or a compound of the formula (1); and




9

5

b) where necessary, conversion of a compound of the formula (9) into a compound of the formula (1).

The compound of the formula (8) in this aspect of the invention may be made by any convenient method such as those described and referenced hereinbefore.

10 Under some conditions for the hydrolysis of the acetal group A in a compound of formula (7), the group R may also simultaneously be hydrolysed which may result in formation of the crystalline lactone (10) [(E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3R,5S)-3,5-dihydroxyhept-6-enoic acid-(3,6)-lactone].



10

Suitable conditions for conversion of compounds of formula (7), for example (7a), into a compound of formula (10) are, for example, treatment with aqueous acid and removal of water by azeotropic distillation of toluene. The crystalline lactone (10) may be isolated instead of the compound (8) and then converted into a compound of the formula (9) or
5 formula (1) by hydrolysis in aqueous acid or base.

In a further aspect of the invention is provided crystalline (*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoic acid-(3,6)-lactone having an X-ray powder diffraction pattern with peaks at 2-theta = 7.9, 15.9 and 20.3.

10 In a further aspect of the invention is provided crystalline (*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoic acid-(3,6)-lactone having an X-ray powder diffraction pattern with peaks at 2-theta = 7.9, 11.9, 15.9, 20.3, 21.7 and 22.5.

15 In a further aspect of the invention is provided crystalline (*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoic acid-(3,6)-lactone having an X-ray powder diffraction pattern substantially as shown in Figure 8.

The utility of the compound of the formula (I) formed by the process of the invention may be demonstrated by standard tests and clinical studies, including those described in EPA
20 521471.

A further aspect of the invention comprises a compound of the formula (I) obtained by the process of the invention as hereinbefore described.

A further aspect of the invention comprises a compound of the formula (I) obtainable by the process of the invention as hereinbefore described.

25 According to a further feature of the invention is a method of treating a disease condition wherein inhibition of HMG CoA reductase is beneficial which comprises administering to a warm-blooded mammal an effective amount of a compound of the formula (I) formed by the process of the invention. The invention also relates to the use of compounds of the formula (I) formed by the process of the invention in the manufacture of a medicament
30 for use in a disease condition.

The compound of the invention may be administered to a warm-blooded animal, particularly a human, in need thereof for treatment of a disease in which HMG CoA reductase is implicated, in the form of a conventional pharmaceutical composition. Therefore in another

aspect of the invention, there is provided a pharmaceutical composition comprising a compound of the formula (I) formed by the process of the invention in admixture with a pharmaceutically acceptable carrier.

Such compositions may be administered in standard manner for the disease condition that it is desired to treat, for example by oral, topical, parenteral, buccal, nasal, vaginal or rectal administration or by inhalation. For these purposes the compound of the formula (I) may be formulated by means known in the art into the form of, for example, tablets, capsules, aqueous or oily solutions, suspensions, emulsions, creams, ointments, gels, nasal sprays, suppositories, finely divided powders or aerosols for inhalation, and for parenteral use (including intravenous, intramuscular or infusion) sterile aqueous or oily solution or suspensions or sterile emulsions. A preferred route of administration is oral. The compound of the formula (I) will be administered to humans at a daily dose in, for example, the ranges set out in EPA 521471. The daily doses may be given in divided doses as necessary, the precise amount received and the route of administration depending on the weight, age and sex of the patient being treated and on the particular disease condition being treated according to principles known in the art.

According to a further feature of the invention, there is provided a process for the manufacture of a pharmaceutical composition containing the compound of the formula (I) as active ingredient, which comprises admixing the compound of the formula (I) together with a pharmaceutically acceptable carrier.

The invention will now be illustrated by the following examples. ¹H NMR were analysed using a Bruker DPX400 operating at a field strength of 400MHz, and unless otherwise stated were run in deuteriochloroform. Chemical shifts are given in parts per million relative to tetramethylsilane.

s = singlet, d = doublet, sept = septet, q = quartet, t = triplet, dd = doublet of doublets, dt = doublet of triplets, m = multiplet

General procedure for synthesis of a compound of formula (7):

Example for R = iso-propyl: iso-Propyl (E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]vinyl}(4R,6S)-2,2-dimethyl[1,3]dioxan-4-yl)acetate

Sodium bis(trimethylsilyl)amide (80.47 mL, 1.0M in tetrahydrofuran (THF)) was added dropwise to a cooled solution of diphenyl [4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-ylmethyl] phosphine oxide (40.43 g, 75 mmol) in THF (477.1 mL) at -65°C over 30 minutes, maintaining the temperature at -65°C . Isopropyl-2-[(4R,6S)-6-formyl-2,2-dimethyl-1,3-dioxan-4-yl] acetate in toluene (21.68 g) was added dropwise to the solution over 35 minutes, maintaining the temperature at -65°C . The contents of the vessel were kept at -65°C for 15 minutes, then allowed to warm evenly to 10°C over 80 minutes. Water (40.4 mL) followed by acetic acid (6.87 g, 114 mmol) were added to give a two phase light yellow solution. The batch was then distilled at atmospheric pressure to remove ~ 485 mL of distillates. This solution was washed sequentially with water (84 mL), 7.0% w/w sodium bicarbonate (92.6 g), 1.8% w/w sodium bicarbonate (91.1 g) and water (63.5 mL). The resulting organic phase was distilled under vacuum at 270 mbar to leave ~ 95 mL of solution in the distillation flask (removing ~ 229 mL of distillates). Methanol (202 mL) at 50°C was charged to the flask and the solution distilled at atmospheric pressure, removing ~ 134 mL of distillates. A further portion of methanol (229 mL) at 50°C was added to the solution and the batch cooled to 40°C over 30 minutes. The batch was cooled to 25°C over 30 minutes, $0-5^{\circ}\text{C}$ over 30 minutes, then chilled to -8°C over 20 minutes and kept at this temperature for 30 minutes. The solid was collected by vacuum filtration, washed with 2 portions of cooled (-8°C) methanol (2 x 80.6 mL) then dried in a vacuum oven at 50°C , 200 mbar, yield = 28.9 g (68.3%).

Analogues with different ester groups R may be made as above using the appropriate starting materials, with the following exceptions:

For the ethyl derivative: after the acetic acid quench the mixture was evaporated to dryness onto silica gel (Merck, 230-400 mesh) and added to a short plug of silica. Elution was performed with 25-27.5% ethyl acetate in iso-hexane. The isolated product was then crystallised from methanol (150 mL) as described in the method above.

For the methyl derivative, the crude product was evaporated to dryness onto silica after the first sodium bicarbonate treatment. Purification was performed on silica gel as for the ethyl derivative (eluting with 14, 16 and 20% ethyl acetate in iso-hexane). The product was crystallised from methanol.

Procedure for compound of formula 7 where R = n-hexyl :

Sodium hydride (141 mg, 60% dispersion in mineral oil, 3.5 mmol) was added in one portion to n-hexanol (15 mL) at ambient temperature. After the resulting effervescence had stopped the clear solution was agitated for 30 minutes. n-Ethyl (E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]vinyl}(4R,6S)-2,2-dimethyl[1,3]dioxan-4-yl)acetate (2 g, 3.6 mmol) was added to the solution in one portion. After 90 minutes, acetic acid (263 mg, 3.6 mmol) was added to the reaction mixture and the solution left at ambient temperature for 16 hours. The solvent was removed on the rotary evaporater (oil pump) and then dissolved in ethyl acetate. The solution was evaporated to dryness onto silica and purified on silica eluting with 20% ethyl acetate in iso-hexane. The resulting pale yellow oil was crystallised from methanol as described above.

Analytical Data: for R-(E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]vinyl}(4R,6S)-2,2-dimethyl[1,3]dioxan-4-yl)acetate

R	¹ H NMR (ppm)
Iso-propyl	1.15 (q, 1H), 1.24 (dd, 6H), 1.27 (dd, 6H), 1.40 (s, 3H), 1.49 (s, 3H), 1.55 (dt, 1H), 2.34 (dd, 1H), 2.50 (dd, 1H), 3.38 (spt, 1H), 3.51 (s, 3H), 3.57 (s, 3H), 4.32 (m, 1H), 4.43 (m, 1H), 5.04 (sptt, 1H), 5.47 (dd, 1H), 6.52 (d, 1H), 7.08 (t, 2H), 7.65 (dd, 2H)
Ethyl	1.14 (q, 1H), 1.25-1.29 (m, 9H), 1.40 (s, 3H), 1.49 (s, 3H), 1.56 (dt, 1H), 2.37 (dd, 1H), 2.55 (dd, 1H), 3.38 (spt, 1H), 3.51 (s, 3H), 3.57 (s, 3H), 4.21-4.12 (m, 2H), 4.37-4.30 (m, 1H), 4.46-4.41 (m, 1H), 5.47 (dd, 1H), 6.53 (d, 1H), 7.08 (t, 2H), 7.65 (dd, 2H)
Methyl	1.14 (q, 1H), 1.27 (dd, 6H), 1.40 (s, 3H), 1.49 (s, 3H), 1.56 (dt, 1H), 2.38 (dd, 1H), 2.57 (dd, 1H), 3.37 (spt, 1H), 3.51 (s, 3H), 3.57 (s, 3H), 3.70 (s, 3H), 4.37-4.30 (m, 1H), 4.46-4.41 (m, 1H), 5.47 (dd, 1H), 6.52 (d, 1H), 7.08 (t, 2H), 7.65 (dd, 2H)

n-Hexyl	0.89 (t, 3H), 1.15 (q, 1H), 1.27 (dd, 6H), 1.37-1.29 (m, 7H), 1.40 (s, 3H), 1.49 (s, 3H), 1.66-1.57 (m, 2H), 2.38 (dd, 1H), 2.55 (dd, 1H), 3.37 (spt, 1H), 3.51 (s, 3H), 3.57 (s, 3H), 4.10 (t, 2H), 4.36-4.29 (m, 1H), 4.46-4.41 (m, 1H), 5.47 (dd, 1H), 6.52 (d, 1H), 7.08 (t, 2H), 7.65 (dd, 2H)
---------	---

General procedure for compounds of formula (8):

5 **Example for iso-propyl analogue:**

Hydrochloric acid (17.24 mL, 0.02M) was added dropwise over 100 minutes to a warm solution of iso-propyl (*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoate (10 g, 17 mmol) in acetonitrile (69 mL) at 35°C. The mixture was agitated for a further 80 minutes then
 10 allowed to cool to 25°C over 30 minutes. Sodium chloride (9.26 g) in water (23 mL) was added to the mixture, which was agitated for 15 minutes then allowed to settle for 16 hours. The organic layer was separated off and the aqueous layer extracted with acetonitrile (15 mL). The combined organic layers were evaporated to dryness and the residue recrystallised from toluene (40 mL). The solid was dried in a vacuum oven at 40°C, 620 mbar, yield = 7.06 g
 15 (79.3%).

Analogues with different ester groups R may be made as above with the following exceptions: for the ethyl derivative, the product does not crystallise from toluene: this solution is evaporated to dryness and purified on silica to afford an oily solid which is then treated with a
 20 1:1 mixture of iso-hexane:toluene and filtered by vacuum filtration.

Analytical Data: for R -(*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoate

R	¹ H NMR (ppm)
t-Butyl	1.27 (d, 6H), 1.42 (dt, 1H), 1.47 (s, 9H), 1.58-1.50 (m, 1H), 2.38 (d, 2H), 3.37 (spt, 1H), 3.51 (s, 3H), 3.57 (s, 3H), 3.65 (bs, 1H), 3.80 (bs, 1H), 4.18-4.15 (m, 1H), 4.48-4.44 (m, 1H), 5.46 (dd, 1H), 6.64 (d, 1H), 7.09 (t, 2H), 7.65 (dd, 2H)

Iso-propyl	1.27-1.25 (m, 12H), 1.45 (dt, 1H), 1.60-1.51 (m, 1H), 2.43 (d, 2H), 3.37 (spt, 1H), 3.51 (s, 3H), 3.57 (s, 3H), 3.57 (bs, 1H), 3.73 (bs, 1H), 4.21-4.19 (m, 1H), 4.47-4.44 (m, 1H), 5.06 (spt, 1H), 5.46 (dd, 1H), 6.64 (d, 1H), 7.09 (t, 2H), 7.65 (dd, 2H)
Ethyl	1.30-1.24 (m, 9H), 1.45 (dt, 1H), 1.60-1.52 (m, 1H), 2.46 (d, 2H), 3.37 (spt, 1H), 3.51 (s, 3H), 3.51 (bs, 1H), 3.57 (s, 3H), 3.70 (bs, 1H), 4.19 (q, 2H), 4.23-4.17 (m, 1H), 4.47-4.45 (m, 1H), 5.46 (dd, 1H), 6.64 (d, 1H), 7.09 (t, 2H), 7.65 (dd, 2H)

Procedure for Formation of Lactone (3R, 5S)

(*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoic acid calcium salt (30.0 g) was dissolved in acetonitrile (300 mL) and saturated brine (50 mL) was added. The solution was then chilled to 0-5°C. The pH was adjusted to 4.0 with a mixture of 4N HCl (15 mL) and saturated brine (35 mL). An extra portion of water (15 mL) was added to dissolve the solid, resulting in two clear phases. The aqueous layer was separated off (112 mL) and the organic phase dried with magnesium sulphate. The solution was distilled at atmospheric pressure until ~100 mL of acetonitrile had been removed, then toluene (250 mL) was added gradually to maintain the distillation flask volume at 200 mL. This resulted in collecting ~390 mL of distillates at a final head temperature of 106°C. The solution was allowed to stir at ambient temperature overnight and was then heated to reflux for two hours. The mixture was cooled to 0-5°C and the resulting solid was filtered, washed with toluene (2 x 20 mL) and dried at 35°C under vacuum.

¹H NMR δ: 1.28-1.26 (m, 6H), 1.69-1.62 (m, 1H), 1.94-1.88 (m, 1H), 2.66-2.60 (m, 1H), 2.72 (dd, 1H), 3.33 (septet, 1H), 3.51 (s, 3H), 3.57 (s, 3H), 4.34-4.30 (m, 1H), 5.26-5.21 (m, 1H), 5.49 (dd, 1H), 6.72 (d, 1H), 7.11 (t, 2H), 7.62 (dd, 2H)

Synthesis of starting materials

Isopropyl- 2-[(4*R*,6*S*)-6-formyl-2,2-dimethyl-1,3-dioxan-4-yl]acetate

Chlorine gas (2469.6 mL, 118 mmol) was charged to toluene (373.3 mL) at -60°C. Dimethyl sulphide (11.67 mL, 121 mmol) was then added dropwise to the cooled solution over 30 minutes, keeping the contents at -60°C. After 30 minutes at this temperature, isopropyl 2-[(4*R*,6*S*)-6-formyl-2,2-dimethyl-1,3-dioxan-4-yl]acetate (24.56 g, 95 mmol) in toluene (46.7

mL) was added dropwise to the vessel over 30 minutes, maintaining the internal temperature at -60°C . The reaction mixture was agitated at -60°C for 30 minutes followed by the dropwise addition of triethylamine (26.36 g, 261 mmol) over 30 minutes, allowing the internal temperature to rise to -50°C . The reaction mixture was then allowed to warm to 25°C evenly over 75 minutes. The resulting slurry was stirred at 25°C for 30 minutes, then water (77 mL) was added and the mixture agitated for 30 minutes. The aqueous layer was separated and the pH checked (pH should be between 7.5 and 8.5). The resulting organic portion was washed with water (23.3 mL) and the organic portion separated for vacuum distillation at 150 mbar. Distillation was continued until ~ 350 mL of toluene had been removed. Toluene (350 mL) was added to the flask and the vacuum distillation repeated at 150 mbar to remove ~ 350 mL of toluene. The resulting solution was transferred to a flask containing activated 4 angstrom molecular sieves and left at ambient temperature overnight. This solution was used directly for the coupling stage.

15 Analogues with different ester groups R may be made as above with the following exceptions: for the methyl analogue, the distillations were performed at much higher vacuum (and therefore at lower temperatures).


Iso-propyl 2-[(4R,6S)-6-formyl-2,2-dimethyl-1,3-dioxan-4-yl]acetate

20 This compound may be made using the procedures described in EP0319847. Analogues with different ester groups R may be made by a similar method.

Diphenyl [4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-ylmethyl] phosphine oxide


25 This compound can be made as described in Patent Application WO00/49014

Figure 1: Methyl (E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]vinyl} (4R,6S)-2,2-dimethyl[1,3]dioxan-4-yl)acetate




2-theta	d-spacing	Relative Intensities
5.7	15.4	51
8.7	10.1	25
9.5	9.3	100
13.6	6.5	54
17.5	5.1	58
19.0	4.7	41
19.9	4.5	53
20.8	4.3	29
21.8	4.1	34
22.4	4.0	41

Figure 2: Ethyl (E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]vinyl}(4R,6S)-2,2-dimethyl[1,3]dioxan-4-yl)acetate



2-theta	d-spacing	Relative Intensities
5.9	15.0	23
8.0	11.0	33
12.2	7.3	21
15.9	5.6	100
18.4	4.8	65
19.5	4.6	82
19.7	4.5	45
23.0	3.9	53
24.3	3.7	37
25.0	3.6	29


Figure 3: iso-Propyl (E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]vinyl}{4R,6S)-2,2-dimethyl[1,3]dioxan-4-yl)acetate



5


2-theta	d-spacing	Relative Intensities
6.0	14.7	35
7.0	12.6	41
7.8	11.3	90
8.7	10.1	37
10.4	8.5	27
11.6	7.6	100
13.0	6.8	20
14.7	6.0	19
15.5	5.7	56
20.2	4.4	53

Figure 4: n-hexyl (E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-methyl(methylsulfonyl)amino]pyrimidin-5-yl}vinyl}{(4R,6S)-2,2-dimethyl[1,3]dioxan-4-yl)acetate




2-theta	d-spacing	Relative Intensities
5.3	16.8	18
7.1	12.4	100
11.4	7.8	10
14.2	6.2	19
14.8	6.0	14
18.9	5.7	27
20.1	4.4	10
20.4	4.4	5
21.4	4.1	12

Figure 5: Ethyl-(*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoate




2-theta	d-spacing	Relative Intensities
4.3	20.4	13
8.1	11.0	100
11.3	7.8	48
12.4	7.1	21
15.1	5.9	18
19.9	4.5	56
21.0	4.2	41
21.7	4.1	18
22.1	4.0	26
23.5	3.8	15

Figure 6: iso-Propyl-(*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoate




2-theta	d-spacing	Relative Intensities
9.8	9.0	88
12.2	7.3	24
13.6	6.5	20
17.3	5.1	57
18.5	4.8	18
19.6	4.5	24
20.1	4.4	53
21.1	4.2	100
22.4	4.0	20
23.3	3.8	16

Figure 7: tert-Butyl-(E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3R,5S)-3,5-dihydroxyhept-6-enoate



2-theta	d-spacing	Relative Intensities
9.8	9.1	31
17.2	5.2	13
18.2	4.9	35
19.4	4.6	15
19.9	4.5	50
20.6	4.3	33
20.8	4.3	100
22.1	4.0	12
25.1	3.5	10
26.3	3.4	18

Figure 8: (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3R,5S)-3,5-dihydroxyhept-6-enoic acid-(3,6)-lactone




5

2-theta	d-spacing	Relative Intensities
7.9	11.1	76
11.9	7.4	25
15.9	5.6	69
20.3	4.4	100
21.7	4.1	59
22.5	3.9	31


Claims

1. A process for the manufacture of the calcium salt of (*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoic acid, comprising:
- 5 a) acid hydrolysis of an acetal protecting group in a compound of the formula (7)



7


wherein A is an acetal or ketal protecting group and R is (1-6C)alkyl, and isolation of the resulting crystalline compound of the formula (8);



8

10


- b) optional recrystallisation of the compound of the formula (8);
- c) hydrolysis of the ester group in the compound (8) to give a dihydroxy carboxylate derivative (9) (wherein M is hydrogen or a metal counterion other than calcium) or a compound of the formula (1); and



9


d) where necessary, conversion of a compound of the formula (9) into a compound of the formula (1).

2. A process as claimed in Claim 1 wherein the the compound of the formula (7) is a compound of the formula (7a).




7a

3. A process for the manufacture of the calcium salt of (*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoic acid, comprising:



8

a) hydrolysis of the ester group in a crystalline compound of the formula (8) (where R is as hereinbefore defined) to give a dihydroxy carboxylate derivative (9) (wherein M is hydrogen or a metal counterion other than calcium) or a compound of the formula (1); and



9

b) where necessary, conversion of a compound of the formula (9) into a compound of the formula (1).

4. A process as claimed in Claim 1 or Claim 2 or Claim 3, wherein R is methyl, ethyl, iso-propyl, tert-butyl or hexyl.
5. A crystalline compound of the formula 7 as defined in Claim 1, which crystalline compound is methyl (E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]vinyl}(4R,6S)-2,2-dimethyl[1,3]dioxan-4-yl)acetate having an X-ray powder diffraction pattern with peaks at 2-theta = 9.5, 13.6 and 17.5.
6. A crystalline compound of the formula 7 as defined in Claim 1, which crystalline compound is ethyl (E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]vinyl}(4R,6S)-2,2-dimethyl[1,3]dioxan-4-yl)acetate having an X-ray powder diffraction pattern with peaks at 2-theta = 15.9, 18.4 and 19.5.
7. A crystalline compound of the formula 7 as defined in Claim 1, which crystalline compound is iso-propyl (E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl]vinyl}(4R,6S)-2,2-dimethyl[1,3]dioxan-4-yl)acetate having an X-ray powder diffraction pattern with peaks at 2-theta = 7.8, 11.6 and 15.5.
8. A crystalline compound of the formula 7 as defined in Claim 1, which crystalline compound is n-hexyl (E)-(6-{2-[4-(4-fluorophenyl)-6-isopropyl-2-

[methyl(methylsulfonyl)amino]pyrimidin-5-yl]vinyl}(4R,6S)-2,2-dimethyl[1,3]dioxan-4-yl)acetate having an X-ray powder diffraction pattern with peaks at 2-theta = 5.3, 7.1 and 18.9.

- 5 9. A crystalline compound of the formula 8 as defined in Claim 1, which crystalline compound is ethyl-(*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoate having an X-ray powder diffraction pattern with peaks at 2-theta = 8.1, 11.3 and 19.9.
- 10 10. A crystalline compound of the formula 8 as defined in Claim 1, which crystalline compound is iso-propyl-(*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoate having an X-ray powder diffraction pattern with peaks at 2-theta = 9.8, 17.3 and 21.1.
- 15 11. A crystalline compound of the formula 8 as defined in Claim 1, which crystalline compound is tert-butyl-(*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoate having an X-ray powder diffraction pattern with peaks at 2-theta = 18.2, 19.9 and 20.8.
- 20 12. Crystalline (*E*)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)amino]pyrimidin-5-yl](3*R*,5*S*)-3,5-dihydroxyhept-6-enoic acid-(3,6)-lactone having an X-ray powder diffraction pattern with peaks at 2-theta = 7.9, 11.9, 15.9, 20.3, 21.7 and 22.5.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.